
SUPPLEMENTAL LABEL

Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1254A-S1 0109

Revus®

Fungicide

For control of certain diseases in listed vegetable crops

Active Ingredient/Guarantee:

Mandipropamid* 23.3%

Other Ingredients: 76.7%

Total: 100.0%

* CAS No. 374726-62-2

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative.

Contains 23.3% Mandipropamid equivalent to 2.08 pounds per gallon or 250 grams per liter of active ingredient

KEEP OUT OF REACH OF CHILDREN.

EPA Reg. No. 1254

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Revus as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Crop	Disease	Rate fl. oz./Acre (lb. a.i./A)	Remarks
Potatoes and other vegetables, tuberous and corm, subgroup <i>See additional crops below.</i>	Late blight <i>(Phytophthora infestans)</i>	5.5 - 8.0 (0.09 - 0.13)	Begin applications prior to disease development and continue throughout the season on a 7-10 day interval. Make no more than 2 consecutive applications before switching to another effective non-Group 40 fungicide. Use the shorter interval and/or higher rates under high pressure or when conditions are conducive to disease.
			The addition of a spreading/penetrating type adjuvant such as a non-ionic surfactant or crop oil concentrate or blend is recommended when applying by ground or air.
Application: For best results, use sufficient water volume to provide thorough coverage. Revus may be applied by ground, chemigation, or aerial application.			

Additional Tuberous and Corm Subgroup Vegetables: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna, Cassava (edible, bitter, and sweet), Chayote (root), Chufa, Dasheen (taro), Ginger, Sweet potato, Taneir, Turmeric, Yam (bean), Yam (true)

Specific Use Restrictions:

- Do not apply more than 32 fl. oz. of product/A/season (0.52 lbs. a.i./A/season).
- Do not apply within 14 days of harvest (14 day PHI).

Crop	Disease	Rate fl. oz./Acre (lb. a.i./A)	Remarks
<p>Tomatoes Tomatillo</p>	<p>Late blight (<i>Phytophthora infestans</i>)</p>	<p>5.5 - 8.0 (0.09 - 0.13)</p>	<p>Begin applications prior to disease development and continue throughout the season on a 7-10 day interval. Make no more than 2 consecutive applications before switching to another effective non-Group 2 fungicide. Use the shorter interval and/or higher rates under high pressure or when conditions are conducive to disease.</p> <p>The addition of a spreading/ penetrating type adjuvant such as a non-ionic surfactant or crop oil concentrate or blend is recommended when applying by ground or air.</p> <p>Application: For best results, use sufficient water volume to provide thorough coverage. Revus may be applied by ground, chemigation, or aerial application.</p>

Specific Use Restrictions:

- Do not apply more than 32 fl. oz. of product/A/season (0.52 lbs. a.i./A/season).
- Do not apply within 1 day of harvest (1 day PHI).

Revus® and the Syngenta logo are trademarks of a Syngenta Group Company

©2009 Syngenta

REV 1254A-S1 0109 – df – 6/25/09