

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ LANNATE® LV INSECTICIDE

EPA Reg. No. 352-384

FOR USE ON FIELD CORN AND SOYBEANS FOR THE CONTROL OF SLUGS IN THE STATES OF DELAWARE, MARYLAND, PENNSYLVANIA, VIRGINIA, AND WEST VIRGINIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

RESTRICTED USE PESTICIDE

DUE TO HIGH ACUTE TOXICITY TO HUMANS

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading or repair or cleaning of application equipment.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ LANNATE® LV can be applied at 1.5 pt/acre when slug populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

APPLICATION

Slugs primarily feed in the early evening, at night or in cool, early morning hours. Slugs are most active when weather conditions are moist and cool. It is recommended to make applications in the early evening through early morning when slugs are active and feeding.

LANNATE® LV should be applied at 5 to 7 day intervals to maintain control. Use sufficient water to obtain thorough, uniform coverage. Since LANNATE® LV is a fast acting contact insecticide, best results follow direct spraying of the target insect.

NOTE:

For Field Corn:

Do not apply LANNATE® LV within 21 days of harvest in field corn.

Do not apply more than 2.25 pounds of methomyl (7.5 pints of LANNATE® LV) per acre per growing season.

Do not make more than a total of 10 applications of LANNATE® LV per crop.

For Soybeans:

Do not apply LANNATE® LV within 14 days of harvest in soybeans.

Do not apply more than 1.35 pounds of methomyl (4.5 pints of LANNATE® LV) per acre per growing season.

Do not make more than a total of 3 applications of LANNATE® LV per crop.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2010 E.I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898