

**Table 1. Average nightly number of *Helicoverpa zea* (corn earworm/bollworm) moths captured in eastern Virginia black light traps, 2017 season.**

Nightly <i>H. zea</i> trap catch average for the week ending:																	
Location (county-town)	6/29	7/06	7/13	7/20	7/27	8/03	8/10	8/17	8/24	8/31	9/07	9/14	9/21	9/28	10/05	10/12	10/19
Chesapeake.....	n/a	n/a	0.0	2.0													
Dinwiddie-Old Hickory .....	n/a	n/a	3.6	5.2													
Dinwiddie-Ford .....	n/a	n/a	n/a	n/a													
Hanover .....	n/a	0.2	2.7	1.1													
Prince George-Templeton .....	n/a	n/a	n/a	0.8													
Prince George-Disputanta .....	n/a	n/a	n/a	0.3													
Richmond Co.-Warsaw .	1.3	0.0	0.0	0.4													
Southampton-Courtland .....	n/a	0.6	0.0	0.0													
Suffolk-Holland.....	n/a	n/a	n/a	8.5													

*n/a = report not available.*

**Table 2. Average nightly number of brown marmorated stink bugs (BMSB) captured in eastern Virginia black light traps, 2017 season.**

Nightly BMSB trap catch average for the week ending:																	
Location (county-town)	6/29	7/06	7/13	7/20	7/27	8/03	8/10	8/17	8/24	8/31	9/07	9/14	9/21	9/28	10/05	10/12	10/19
Chesapeake.....	n/a	n/a	0.0	0.0													
Dinwiddie-Old Hickory .....	n/a	n/a	0.0	0.0													
Dinwiddie-Ford .....	n/a	n/a	n/a	n/a													
Hanover .....	n/a	1.0	1.9	2.7													
Prince George-Templeton .....	n/a	n/a	n/a	0.2													
Prince George-Disputanta .....	n/a	n/a	n/a	0.0													
Richmond Co.-Warsaw .	0.0	0.0	0.1	1.4													
Southampton-Courtland .....	n/a	0.0	0.0	0.0													
Suffolk-Holland.....	n/a	n/a	n/a	0.0													

*n/a = report not available.*