

Supplemental Labeling for Cucumbers



SANDEA[®] is a selective herbicide for control of listed broadleaf weeds and nutsedge

ACTIVE INGREDIENT:	% BY WT.
Halosulfuron-methyl, methyl 3-chloro-5-(4,6-dimethoxypyrimidin-2-ylcarbamoylsulfamoyl)	75.0%
-1-methylpyrazole-4-carboxylate.....	25.0%
OTHER INGREDIENTS	TOTAL 100.0%

EPA Reg. No. 81880-18-10163

This label expires 05-31-2018 and must not be used or distributed after that date.

KEEP OUT OF REACH OF CHILDREN CAUTION

Read the label affixed to the container for Sandea Herbicide before applying.

- The labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- Use of Sandea herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Sandea Herbicide.

CROP	OZ/ACRE	DIRECTIONS FOR USE
CUCUMBERS (14) <i>(including pickles)</i> CANTALOUPE (57), HONEYDEWS (57), AND CRENSHAW MELONS (57)	1/2 - 1	<p>Apply uniformly with ground equipment in a minimum of 15 gal of water per acre.</p> <p>Direct-seeded: Bare ground (no mulch)</p> <ul style="list-style-type: none"> • Preemergence - Apply SANDEA after planting, but prior to soil cracking. Use the lower rate on lighter textured soils with low organic matter. • Postemergence - Apply SANDEA after the crop has reached at least 3 to 5 true leaves but before first female flowers appear. SANDEA can be applied as an over-the-top application, a directed spray application, or with crop shields to minimize contact of the herbicide with the crop. <p>Direct-seeded: Plastic mulch</p> <ul style="list-style-type: none"> • Pre-seeding - Apply SANDEA following final bed shaping and just prior to the installation of the plastic mulch. Crop may be seeded into this treated area no sooner than 7 days after application and the installation of the plastic mulch unless local conditions demonstrate safety at an earlier interval. Use the lower rate on lighter textured soils with low organic matter. • Postemergence - Apply SANDEA after the crop has at least 3 to 5 true leaves but before first female flowers appear. SANDEA can be applied as an over-the-top application, a directed spray application, or with crop shields to minimize contact of the herbicide with the crop. Additional phytotoxicity may occur when applications are made over plastic due to concentration of product in the planting hole. Note: Over-the-top applications on plastic are not allowed in Northeastern and Midwestern states. <p>Transplanted: Bare ground (no mulch)</p> <ul style="list-style-type: none"> • Pre-transplant - Apply SANDEA as a pre-transplant application. Crop may be transplanted into this treated area no sooner than 7 days after application unless local conditions demonstrate safety at an earlier interval. Use the lower rate on lighter textured soils with low organic matter. Care should be taken to limit movement of SANDEA-treated surface soil during the transplanting process since if treated soils is moved into the transplant hole injury can occur. • Post-transplant - Apply SANDEA to transplants that are established and actively growing. Applications should not be made until plants are actively growing and in the 3 to 5 true leaf stage or no sooner than 14 days after transplanting unless local conditions demonstrate safety at an earlier interval, but before first female flowers appear. SANDEA may be applied as an over-the-top application, a directed spray application, or with crop shields to minimize contact of the herbicide with the crop. <p>Transplanted: Plastic mulch</p> <ul style="list-style-type: none"> • Pre-transplant - Apply SANDEA following final bed shaping and just prior to the installation of the plastic mulch. Crop may be transplanted into this treated area no sooner than 7 days after the application and the installation of the plastic mulch unless local conditions demonstrate safety at an earlier interval. Use the lower rate on lighter textured soils with low organic matter. Care should be taken to limit movement of SANDEA-treated surface soil during the transplanting process since if treated soils is moved into the transplant hole injury can occur. • Post-transplant - Apply SANDEA to transplants that are established, actively growing and in the 3 to 5 true leaf stage or no sooner than 14 days after transplanting unless local conditions demonstrate safety at an earlier interval, but before first female flowers appear. Apply SANDEA as an over-the-top application, a directed spray application, or with crop shields to minimize contact of the herbicide with the crop. Additional phytotoxicity can occur when applications are made over plastic due to concentration of product in the transplant hole. Note: Over-the-top applications on plastic are not allowed in Northeastern and Midwestern states.

CROP	OZ/ACRE	DIRECTIONS FOR USE
		<p>Direct-seeded and Transplant:</p> <ul style="list-style-type: none"> • Row Middle/Furrow Applications - Apply SANDEA between rows of direct-seeded or transplanted crop. Avoid contact of the herbicide with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Reduce rate and spray volume in proportion to area actually sprayed. <p>Split Applications for Nutsedge:</p> <ul style="list-style-type: none"> • Preemergence followed by postemergence for nutsedge control To maximize control of nutsedge, it may be necessary to use a postemergence application to those areas where the nutsedge has emerged later following a preemergence application. For these situations, use a spot treatment method treating only those areas of emerged nutsedge. Application rate should not exceed 1.0 oz product per treated acre in these areas. Use a water volume that will allow for good coverage of the plants. Avoid contact of the herbicide with the planted crop. • Postemergence followed by postemergence for nutsedge control To maximize control of nutsedge, it may be necessary to use a second postemergence spot application to those areas where the nutsedge has emerged or re-grown. For these situations, use a spot treatment method treating only those areas of emerged nutsedge. Allow a minimum of 21 days between applications. Application rate should not exceed 1.0 oz product per treated acre in these areas. Use a water volume that will allow for good coverage of the plants. Avoid contact of the herbicide with the planted crop. <p>PRECAUTIONS:</p> <ul style="list-style-type: none"> • Runners that come in contact with the plastic can pick up residual SANDEA and may exhibit a visual crop response. • Consult "Use Precautions" and "For Optimum Results" for important usage information. <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • Do not apply more than 2 applications or 2 oz/A of product by weight (0.094 lb a.i./acre) per 12 month period. (includes applications to the crop and to row middle/furrows)

Supplemental label Cucumbers (EPA approved 5-11-16)



EPA Reg. No. 81880-18-10163
EPA Est. No.

Produced For:
Canyon Group LLC
C/O Gowan Company
PO Box 5569
Yuma, AZ 85366-5569